



УДК 504.4.062

## ALGORITHM FOR DETERMINING OF ENVIRONMENTAL SAFETY OF DRINKING WATER SUPPLY

Krysinska D. O. / Крисінська Д. О.

*Lecturer of Department of Ecology/ викладач кафедри екології*

ORCID: 0000-0002-3117-6039

*Petro Mohyla Black Sea National University, Mykolaiv, 68 Desantnykiv, 10, 54000**Чорноморський національний університет імені Петра Могили,**Миколаїв, вул. 68 Десантників, 10, 54000*

**Abstract.** *An urgent practical task, the study of approaches to determining the environmental safety of drinking water supply, is considered in the article. The author has analyzed various Ukrainian and international methodological approaches to determining the environmental safety of drinking water supply. It has been found that the most effective is the use of environmental risk assessment, which appears to be the main tool for environmental safety assessment. On the basis of the analyzed methods, the complex one, which is expressed in the form of the algorithm for determining the environmental safety of drinking water supply, has been created.*

**Key words:** *environmental safety, drinking water supply, environmental risk, drinking water, industrial water, dual systems, assessment algorithm.*

**Introduction** Building algorithms for assessing the impact of a negative factor on the environment or its components is a common practice in environmental, technical, mathematical research, as it allows to identify, classify and structure factors, determine the level of risk manifestation of negative consequences, and therefore assess the level of danger and make necessary decisions.

The advantages of using algorithms for assessing environmental safety are reflected in the works [1-7]. In these works, algorithms are presented in the form of block diagrams for assessing environmental safety and risks of the impact of various chemicals on ecosystems and their subsystems, as well as living organisms that live in this environment and are directly affected.

Taking into account the experience of previous scientists, we propose to assess the environmental safety of drinking water supply using algorithms.

It is worth noting that the ecological safety of drinking water supply (ESDWS) is understood as a state of drinking water supply, in which the safety ranges of the risk of negative effects are established, while minimizing adverse effects on ecosystem components, primarily on humans, provided that the necessary scientifically reasonable economic and energy costs are used.

**Main part** In our study, the determination of the environmental safety of drinking water supply was carried out through calculations of the environmental risk as the main tool for assessing environmental safety in general, using methods [8,9], combined into a comprehensive approach that takes into account carcinogenic and non-carcinogenic effects, where the range of safety of factor effect is ultimately determined.

The goal of environmental safety is to reduce morbidity, mortality, to increase the duration and quality of human life. The quality of drinking water is one of the main factors of impact on a human in the systems of environmental safety of drinking water supply. It depends on the initial quality of water of the water supply source,



water treatment technology, technical condition of water supply networks, etc.

Absolute achievement of this goal is possible with the constant control and monitoring of sources of influence and factors which disrupt the state of ecological systems. In the case of environmental safety of drinking water supply, the first stage of control should be carried out by the state and organizations responsible for collecting data on the state of water bodies that are sources of water supply. At the same time, it is important to continuously monitor those enterprises that discharge wastewater and to introduce new approaches to water intake, which will consist in permitting its implementation, downstream of the discharge site. Thus, there will be a natural process of stimulating enterprises to improve the quality of treatment of wastewater, which, when mixed with water, will immediately become a new resource for production.

From the point of impact of the level of environmental safety of drinking water supply on a human, it is most effective to conduct the assessment through the quality of drinking water, but the ACL coefficient and methodological approach to comparing the values of monitoring data and standard are considered to be ineffective due to the lack of understanding of the impact mechanism, the manifestation of harm in case of exceeding the standardized value. The most effective tool, which at this stage of development of scientific research as for the assessment of the impact of anthropogenic activities on the human body, is environmental risk. Figures 1-3 show the algorithms for determining the environmental safety of drinking water supply - block diagram 1 and environmental risk assessment - block diagram 2 for carcinogenic effects, block diagram 3 for non-carcinogenic.

The block diagram 1 (fig. 1) is a universal tool for assessing ESDWS. At the first stages of the assessment, after the establishment of the affecting factors on the object of study, the collection and analysis of monitoring and statistical data begins. After the formation of the database, the intermediate indicators of the average annual, average daily doses of the values, which are necessary for further calculations, are determined.


One of the main stages of the algorithm is the stage "Determination of the environmental risk of carcinogenic and non-carcinogenic", the calculations are performed using block diagram 2 (fig. 2) [8] and block diagram 3 (fig. 3) [9], which are mandatory components of the assessment of ESDWS. After establishing the values of environmental risks, if necessary, modeling and/or forecasting is carried out, which will allow to predict a change in the trend of the indicator in the future, or to create a statistical series, in the case of absence of real monitoring data.

In the process of establishing the risk limit, the range of impact safety for indicators, which will be determined by the limits, is determined:

$$0.00001 \leq R_c \leq 0.00001 - \text{for carcinogenic impact factors;}$$

$$0.16 \leq R_{nc} \leq 0.02 - \text{for non-carcinogenic impact factors.}$$


The novelty of the proposed method of determining the level of environmental safety of drinking water supply is that as a result of the assessment we will obtain not just a number which will show the excess of the ACL standard for a particular substance in drinking water, but a complex value that will characterize the impact on a human.




**Fig. 1. Algorithm for determining the environmental safety of drinking water supply**

**Ending and conclusions**

The final step towards establishing the level of ESDWS is the assessing of the risk limits, namely, within the range - "safely", outside the range "dangerously". To reduce conceptual confusion, for clarity and intelligibility of wording, we suggest using only two levels of safety assessment - "safe" and "dangerous" for assessing ESDWS. In the case of a safe assessment result, the system and its components are in a state of sustainable development. In the case of a hazardous one, it is necessary to take a number of urgent measures, ranging from determining the nature of the consequences of the hazard, making decisions on replacing technologies, management and economic costs, completing with reassessing the risks until a "safe" result is achieved.



**Fig. 2. Algorithm for determining potential carcinogenic risk**



**Fig. 3. Algorithm for determining potential non-carcinogenic risk**

**References:**

1. Akimov V. A. Fundamentals of risk analysis and management in natural and man-made spheres/ V. A. Akimov, V. V. Lesnyh, N. N. Radaev. – M.: Business express, 2004. – 352 p.
2. Voitenko L.V. The concept of integrated water quality assessment for different types of water consumption using the Harrington desirability function/ Voitenko L.V., Kopilevich V. A., Strokal M. P. – K.: Bioresources and nature



management: chemistry. – Volume 7, # 1-2, 2015. – P. 25-36. – [Electronic resource] - Access mode: [http://irbis-nbuv.gov.ua/cgi-bin/irbis\\_nbuv/cgiirbis\\_64.exe?C21COM=2&I21DBN=UJRN&P21DBN=UJRN&IMAGE\\_FILE\\_DOWNLOAD=1&Image\\_file\\_name=PDF/Zp\\_2019\\_1\\_5.pdf](http://irbis-nbuv.gov.ua/cgi-bin/irbis_nbuv/cgiirbis_64.exe?C21COM=2&I21DBN=UJRN&P21DBN=UJRN&IMAGE_FILE_DOWNLOAD=1&Image_file_name=PDF/Zp_2019_1_5.pdf)

3. Bardina D. A. Development of an algorithm for assessing the risk to public health when exposed to chemicals that pollute drinking water/ D. A. Bardina, P. G. Mikhailova. – Advances in chemistry and chemical technology. Volume XXIX, Moscow: 2015, # 4 – P. 57-59. – [Electronic resource] - Access mode: <https://cyberleninka.ru/article/v/razrabotka-algoritma-otsenki-riska-zdorovyu-naseleniya-pri-vozdeystvii-himicheskikh-veschestv-zagryaznyayuschih-pitievuyu-vodu>

4. Bezsonov, Ye. M. Determining the level of ecological safety of the region by the method of toxic-energy response of biotic components of aquatic ecosystems: The dissertation on competition of a scientific degree of the candidate of technical sciences on a specialty 21.06.01 "Ecological safety". Lviv: Natsionalnyi universytet «Lvivska politekhnik», 2018.

5. Dobrovolsky V. V. Ecological risk: assessment and management: textbook [text] / V. V. Dobrovolsky. – Mykolaiv: PMBSSU, 2010. – 216 p.

6. Krysinska D. O. Algorithm for determining environmental risk in the system of ecological safety of drinking water supply/ D. O. Krysinska, N. O. Voskoboynikova // Natural water resources of the Carpathian region. Problems of protection and rational use. Proceedings of the Thirteenth International Scientific and Practical Conference: a collection of scientific articles. – Lviv, 2014. – P. 29-32.

7. Orel S. M., Malovanyy M. S., Orel D. S. Assessment of ecological risk. Impact on human health [textbook]. – Lviv, publishing house LPNU, 2013. – 224 p.

8. Order of the Ministry of Health of Ukraine of October 21, 2005 N 545 On approval of guidelines "Assessment of carcinogenic risk to public health from the consumption of chlorinated drinking water" – [Electronic resource] - Access mode: <http://mozdocs.kiev.ua/view.php?id=4448>.

9. Methodical recommendations of 05/30/1997 RF No. 2510 / 5716-97-32 "Comprehensive hygienic assessment of the degree of tension in the medical and ecological situation in various territories, caused by toxicant pollution of the population's environment" – [Electronic resource] - Access mode: [http://www.lawrussia.ru/texts/legal\\_744/doc\\_744a\\_498x\\_422.html](http://www.lawrussia.ru/texts/legal_744/doc_744a_498x_422.html).

**Анотація.** В статті розглядається актуальна практична задача – вивчення підходів до визначення екологічної безпеки питного водопостачання. Автором проаналізовано різні українські і міжнародні методичні підходи до визначення екологічної безпеки питного водопостачання. Встановлено, що найбільш ефективним є використання оцінки екологічного ризику, який постає головним інструментом оцінки екологічної безпеки. На основі проаналізованих методик створено комплексну, яка виражена у вигляді алгоритму визначення екологічної безпеки питного водопостачання.

**Ключові слова:** екологічна безпека, питне водопостачання, екологічний ризик, питна вода, технічна вода, дуальні системи, алгоритм оцінювання.



*Стаття підготовлена в рамках виконання науково-дослідної теми  
«Підвищення рівня екологічної безпеки питного водозабезпечення в  
Миколаївській області» (№ ДР 0112U005117) кафедри екології  
Чорноморського національного університету імені Петра Могили.*

*Науковий керівник: д.т.н., проф. Клименко Л. П.*

© Крисінська Д. О.